

VETMEC PALATABLE ALLWORMER FOR DOGS IS

to be used as a single dose treatment. Tablets are palatable and may be administered directly as a treat or crushed and put on food. Fleas are intermediate hosts for tapeworm (*Dipylidium caninum*). Effective flea control will reduce the incidence of tapeworm.

SAFETY DIRECTIONS: May irritate the eyes. Avoid contact with eyes. Wash hands after use.

FIRST AID: If poisoning occurs, contact a doctor or Poisons Information Centre. Telephone Australia 131 126.

STORAGE: Store below 30°C (Room Temperature).

DISPOSAL: Dispose of empty container by wrapping with paper and putting in garbage.

APVMA: 64703/10/0210

1P0569

[B]: EXP:



CHEMVET AUSTRALIA PTY. LIMITED
1/8 ROCKLEA DRIVE,
PORT MELBOURNE 3207
1800 243 683 www.chemvet.com.au



CAUTION
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS
FOR ANIMAL TREATMENT ONLY

Vetmec Palatable Allwormer Tablets For Dogs



ACTIVE CONSTITUENTS: Each tablet contains:

- 100 mg PRAZIQUANTEL
- 450 mg OXIBENDAZOLE

FOR THE TREATMENT AND CONTROL OF:

Roundworm • Hookworm • Whipworm
• Tapeworm (*Dipylidium caninum*)
including hydatids (*Echinococcus granulosus*) in dogs. Does not control heartworm in dogs.

10 TABLETS

1 Tablet Per 20kg Bodyweight

For use against the following species of worms:

Toxascaris leonina, *Toxocara canis*, *Ancylostoma caninum*,
Uncinaria stenocephala, *Trichuris vulpis*, *Dipylidium caninum*,
Taenia ovis, *Taenia pisiformis*, *Taenia hydatigena* and
Echinococcus granulosus (hydatids) in dogs.

DIRECTIONS FOR USE:

DOSAGE AND ADMINISTRATION:

1 tablet/20kg bodyweight

PUPPIES: Treat at 2, 4, 8 and 12 weeks of age.

ADULTS: Treat every 3 months.

BREEDING BITCH: Treat during gestation, one week before birth and 3 to 4 weeks after birth.

Dogs should not be fed offal from any species. Dogs in hydatid areas should be treated every 6 weeks. Where whipworm is a problem, treat every 6-8 weeks after 3 months of age. If worm problems persist, consult a veterinarian.

