

64347_48129_10L_immediate_MPL_V02.pdf

800mm x 150mm

For A3 print @ 47% For A4 print at @ 35%

This comment box is not part of the label

VETMEC DUAL

ORAL DRENCH FOR SHEEP AND LAMBS

DIRECTION FOR USE

RESTRAINTS

DO NOT USE on sheep which are producing or may in the future produce milk or milk products for human consumption.

PRECAUTIONS

Exercise care in handling weak, pregnant and young animals to avoid unnecessary stress.

Avoid yarding animals off-feed overnight and ensure animals have access to water when yarded prior to drenching. The recommended dose should not be exceeded except under veterinary supervision.

Maintain drench gun carefully to ensure accurate dosing. Before and after use, clean the drenching gun by flushing out with cold water. DO NOT use hot water.

SHAKE CONTAINER BEFORE USE

EFFICIECY SPECTRUM (MATURE & IMMATURE)

Including benzimidazole or levamisole resistant worms. This product is not effective against worms that are resistant to BOTH benzimidazole and levamisole compounds)

ROUND WORMS

Barbers Pole Worm	(<i>Haemonchus contortus</i>)
Small Brown Stomach Worms	(<i>Ostertagia spp.</i>)
Black Scour Worms	(<i>Trichostrongylus spp.</i>)
Stomach Hair Worm	(<i>Trichostrongylus axei</i>)
Small Intestinal Worms	(<i>Cooperia spp.</i>)
Thin Necked Intestinal Worms	(<i>Nematodirus spp.</i>)
Hookworms	(<i>Bunostomum spp.</i>)
Nodule Worm	(<i>Oesophagostomum columbianum</i>)
Large Bowel Worm	(<i>Oesophagostomum venulosum</i>)
Large Mouth Bowel Worm	(<i>Chabertia ovina</i>)

LUNGWORM

Large Lung Worm (*Dictyocaulus filaria*)

TAPEWORM

Heads and Segments (*Moniezia spp.*)

OVICIDAL

Roundworm and Lungworm eggs.

Resistance may develop to any drench. Ask your local veterinary practitioner or animal health advisor for recommended parasite management practises for your area to reduce development of resistance. It is advisable that a drench-resistance test be conducted before any drench is used.

DOSAGE AND ADMINISTRATION

1mL Vetmec Dual per 10 Kg bodyweight.

Drench sheep orally, using an appropriate drench gun.

During drenching, regularly check drench gun to ensure it is delivering the correct dose.

A representative sample of the animals should be weighed before treatment.

Dose the mob to the heaviest animal by liveweight in each group (ewes, wethers, rams, lambs). Do not underdose.

Where there is a large variation in size within the group, dose rate should be based on the label directions for each weight range. Drafting into two lines may be appropriate, to avoid excessive overdosing.

VETMEC DUAL

ORAL DRENCH FOR SHEEP AND LAMBS

VETMEC DUAL ORAL DRENCH FOR SHEEP AND LAMBS

Bodyweight Kg	Dose Volume mL	10L Pack Treats
Up to 20	2.0	5000
21 - 25	2.5	4000
26 - 30	3.0	3333
31 - 35	3.5	2857
36 - 40	4.0	2500
41 - 45	4.5	2222
46 - 50	5.0	2000
51 - 55	5.5	1818
56 - 60	6.0	1666
61 - 65	6.5	1538
66 - 70	7.0	1428
71 - 75	7.5	1333

Sheep in excess of 75 kg should be dosed at 1 ml/10 kg

WITHHOLDING PERIODS:

MEAT: DO NOT USE LESS THAN 10 DAYS BEFORE SLAUGHTER FOR HUMAN CONSUMPTION.
MILK: DO NOT USE ON SHEEP WHICH ARE PRODUCING OR MAY IN THE FUTURE PRODUCE MILK OR MILK PRODUCTS FOR HUMAN CONSUMPTION.

TRADE ADVICE

EXPORT SLAUGHTER INTERVAL (ESI): DO NOT USE less than 10 days before slaughter for export. The ESI on this label was correct at the time of label approval. Before using this product, confirm the correct ESI from Chemvet Australia on 1800 243683 or the APVMA website www.apvma.gov.au/residues/ESI.shtml

SAFETY DIRECTIONS

Poisonous if swallowed. Harmful if absorbed by skin contact. Avoid contact with eyes and skin. Wear rubber gloves. Wash hands after use.

FIRST AID

If poisoning occurs contact a doctor or the Poisons Information Centre. (Ph:131 126).

Additional information is available in the Material Safety Data Sheet for this product. For a copy phone 1800 243 683.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND THE ENVIRONMENT

Albendazole and Levamisole are extremely toxic to aquatic species. DO NOT contaminate dams, rivers or streams or other water ways with the drench or used containers.

DISPOSAL

This container can be recycled if it is clean, dry, free of viable residues and has the *DrumMUSTER* logo visible. Triple or pressure rinse container for disposal. Dispose of rinsate or any undiluted material according to State legislative requirements. Wash outside of the container and the cap. Store cleaned container in a sheltered place with cap removed. It will then be acceptable for recycling at any *DrumMuster* collection or similar container management program site. The cap should not be replaced but may be taken separately.

STORAGE

Store below 30°C (room temperature) in original container, tightly closed in a dry place.

MANUFACTURERS WARRANTY & EXCLUSION OF LIABILITY

This product is warranted fit for the purposes specifically recommended by Chemvet Australia Pty Ltd when used strictly as directed on this label. All other warranties and obligations or liabilities whether expressed or implied by statute or otherwise, are excluded to the full extent that exclusion is permitted by law.

Chemvet Australia Pty Ltd. 1/8 Rocklea Drive, Port Melbourne, VIC 3207 Ph: 1800 243 683

CAUTION

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING
FOR ANIMAL TREATMENT ONLY

VETMEC DUAL

ORAL DRENCH FOR SHEEP AND LAMBS

Albendazole Oxide 36.0 g/L (Equivalent to 34.0 g/L Albendazole)
Levamisole Hydrochloride 82.5 g/L (Equivalent to 70.0 g/L Levamisole)



For the control of gastrointestinal roundworms sensitive to benzimidazoles and levamisole including those resistant to either benzimidazoles or levamisole. For control of lungworm and tapeworm. Reduces the output of viable worm eggs.

10
LITRES

SHAKE WELL BEFORE USE



Chemvet Australia Pty Ltd.
1/8 Rocklea Drive, Port Melbourne, VIC 3207
Ph: 1800 243 683
Website: www.chemvet.com.au