



Product Name: VETMEC LEV POUR-ON FOR CATTLE  
APVMA Approval No: 64597/120378



Label Name:	VETMEC LEV POUR-ON FOR CATTLE
-------------	-------------------------------

Signal Headings:	POISON KEEP OUT OF REACH OF CHILDREN FOR ANIMAL TREATMENT ONLY READ SAFETY DIRECTIONS BEFORE OPENING OR USING
------------------	--

Constituent Statements:	200 g/L LEVAMISOLE (AS LEVAMISOLE BASE) SOLVENT: 150 g/L Diethylene Glycol Monobutyl Ether
-------------------------	---

Claims:	<p>For the treatment of levamisole sensitive gastro-intestinal parasites and lungworm in cattle</p> <p>EFFICACY SPECTRUM MATURE AND IMMATURE</p> <p>VETMEC LEV POUR ON FOR CATTLE contains levamisole, a member of the imidazothiazole family of drenches and is effective against sensitive strains of the following parasites:</p> <p>Ostertagia ostertagi (Small brown stomach worm (mature adults only) in cattle under 12 months of age*) Haemonchus placil (Barber's pole worm) Trichostrongylus axei (Stomach hair worm) Trichostrongylus.spp (Black scour worm) Bunostomum phlebotomum (Hook worm) Oesophagostomum radiatum (Nodule worm) Cooperia spp. (Small intestinal worm) Nematodirus spp. (Thin necked intestinal worm) Dictyocaulus viviparous (Lungworm)</p> <p>*Where Ostertagia is a problem in cattle over 12 months old, products containing benzimidazoles or avermectins/milbemycins are recommended</p> <p>Resistance may develop to any drench</p>
---------	---

Net Contents:	10L 15L 1L 2.5L 20L 25L
---------------	--

5L

Directions for Use:

Restraints:

Do not use in lactating cows where milk may be used or processed for human consumption  
**FOR USE ON CATTLE ONLY**  
Do not treat wet animals  
Do not use on dogs or horses  
Do not use this product as an oral drench or as an injection

Contraindications:

Precautions:

Exercise care in handling weak, pregnant and young animals to avoid unnecessary stress. Avoid yarding animals off-feed overnight and ensure animals have access to water when yarded prior to drenching. Recommended dose should not be exceeded, except under veterinary supervision.

Overdose may cause animals to show temporary excitement, salivation or muscular tremors.

Side Effects:

Dosage and Administration:

This section contains file attachment.

General Directions:

VETMEC L POUR ON FOR CATTLE may be used concurrently with pour-on products for lice control provided the products are applied to different parts of the backline.

Withholding Periods:

MEAT: DO NOT USE less than 3 days before slaughter for human consumption.  
MILK: DO NOT USE in lactating cows where milk may be used or processed for human consumption

Trade Advice:

EXPORT SLAUGHTER INTERVAL (ESI): DO NOT USE less than 7 days before slaughter for export.

The ESI on this label was correct at the time of label approval. Before using this product, confirm the current ESI from Chemvet Australia Pty Ltd on 1800243683 or the APVMA website ([www.apvma.gov.au](http://www.apvma.gov.au)).

Safety Directions:

Poisonous if swallowed. May irritate the eyes and skin. Avoid contact with eyes and skin and wear rubber gloves

First Aid Instructions:

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone 131126.  
If skin contact occurs, remove contaminated clothing and wash skin thoroughly. If in eyes, hold eyes open, flood with water for at least 15 minutes and see a doctor.

First Aid Warnings:	
---------------------	--

Additional User Safety:	Additional information is listed in the Safety Data Sheet.
-------------------------	--

Environmental Statements:	<p>Environmental protection:</p> <p>Do not contaminate dams, rivers or streams with the drench or used container.</p>
---------------------------	---

Disposal:	<p>(1L): Dispose of container by wrapping with paper and putting in garbage.</p> <p>(2.5L, 5L, 10L, 15L, 20L, 25): Triple or( preferably) pressure rinse container. DO NOT dispose of undiluted chemicals on-site. If recycling, replace cap and return clean container to recycler or designated collection point. If not recycling, break, crush, or puncture container and deliver to an approved waste management facility. If an approved waste management facility is not available, bury the broken, crushed or punctured containers 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.</p> <p>If the container has the drumMUSTER logo visible, and has been thoroughly cleaned and dried, and is free of any visible residues, it can be recycled at any drumMUSTER collection or similar container management program site. In this cast the cap should not be replaced, but may be recycled separately with the container.</p>
-----------	---

Storage:	Store below 30 °C (room temperature) in tightly closed original container. Protect from light, and store in a well ventilated dry place
----------	---

## DOSAGE AND ADMINISTRATION

Calibrate applicator gun before use to ensure accurate dosage. After use, clean gun by flushing out with clean cold water. Do not use hot water.

### DOSE RATE

Apply 2.5mL per 50kg bodyweight evenly along the backline using the appropriate applicator gun (equivalent to 10mg of Levamisole base per kg liveweight). For accurate application, the nozzle should be in contact with the skin.

Liveweight (Kg)	Dose (mL)
Up to 50 Kg	2.5
51-100	5.0
101-150	7.5
151-200	10.0
201-250	12.5
251-300	15.0
301-350	17.5
351-400	20.0
401-450	22.5
451-500	25.0
501-550	27.5
551-600	30.0
601-650	32.5

Always dose on a bodyweight basis.

A representative sample of animals should be weighed before treatment either with scales or a weighband. Dose rate should be based on the heaviest cattle in each group (cows, steers, calves, bulls).

DO NOT underdose.

Where there is a large variation in size within the group, draft into two or more lines based on bodyweight, to avoid excessive overdosing.