

Product Name: VETMEC POUR-ON FOR CATTLE  
APVMA Approval No: 84128/118387



Label Name:	VETMEC POUR-ON FOR CATTLE
-------------	---------------------------

Signal Headings:	CAUTION KEEP OUT OF REACH OF CHILDREN FOR ANIMAL TREATMENT ONLY READ SAFETY DIRECTIONS BEFORE OPENING OR USING
------------------	---

Constituent Statements:	10 g/L IVERMECTIN SOLVENT: 150 g/L Diethylene glycol monobutyl ether
-------------------------	---

Claims:	<p>FOR THE TREATMENT AND CONTROL OF IVERMECTIN SENSITIVE INTERNAL AND EXTERNAL PARASITES OF BEEF AND DAIRY CATTLE.</p> <p>INDICATIONS: For the treatment and control of adult and some immature stages of: Gastro-intestinal roundworms:</p> <p>Barbers Pole Worm (<i>Haemonchus placei</i>) Hookworm (<i>Bunostomum phlebotomum</i>) Intestinal Threadworm (<i>Strongyloides papillosus</i>) Large Bowel Worm (<i>Oesophagostomum radiatum</i>) Small Intestinal Worm (<i>Cooperia oncophora</i>) (<i>Cooperia punctata</i>) Small Brown Stomach Worm (<i>Ostertagia ostertagi</i>) (including inhibited stages) Stomach Hair Worm (<i>Trichostrongylus axei</i>) Thin Necked Intestinal Worm (<i>Nematodirus spathiger</i>) Whipworm (<i>Trichuris</i> spp.) Lungworm (<i>Dictyocaulus viviparus</i>)</p> <p>Sucking and Biting lice: Biting lice (<i>Bovicola bovis</i>) Little sucking lice (<i>Solenopotes capillatus</i>) Long nose sucking lice (<i>Linognathus vituli</i>) Short nosed sucking lice (<i>Haematopinus eurysternus</i>)</p> <p>Mites (<i>Chorioptes bovis</i>, <i>Sarcoptes scabiei</i> var <i>bovis</i>)</p> <p>Buffalo Fly (<i>Haematobia irritans exigua</i>)</p> <p>Cattle Tick (<i>Rhipicephalus (Boophilus) microplus</i>)</p> <p>VETMEC POUR-ON FOR CATTLE contains ivermectin, which has a broad spectrum of activity against most internal and external parasites with sensitivity to ivermectin. Ivermectin is a member of the macrocyclic lactone anthelmintic family of chemicals. Resistance can develop to any chemical.</p>
---------	--

	<p>Maximum tick efficacy is achieved 4-5 days after treatment. In planning a strategic tick control program, it should be noted that viable eggs may have been laid by engorged female ticks that drop from cattle in the initial days after treatment. VETMEC POUR-ON FOR CATTLE should be alternated with conventional dips or Pour-Ons when the program requires repeat treatments.</p> <p>SUSTAINED ACTIVITY: VETMEC POUR-ON FOR CATTLE applied at the recommended dose of 1mL per 20kg bodyweight effectively controls infection with <i>Ostertagia</i> spp. (Small brown stomach worm) acquired up to 14 days after treatment, with <i>Dictyocaulus viviparus</i> (Lungworm) acquired up to 28 days after treatment and <i>Haematobia irritans exigua</i> (Buffalo Fly) up to 28 days after treatment.</p>
--	--

Net Contents:	500mL, 1L, 2.5L, 10L, 12.5L, 15L, 20L, and 25L
---------------	--

Directions for Use:	
---------------------	--

Restrains:	Retreatment interval: DO NOT retreat animals for 42 days after last treatment.
------------	--

Contraindications:	
--------------------	--

Precautions:	
--------------	--

Side Effects:	Temporary hair loss occasionally occurs in the application area but will resolve without further treatment.
---------------	---

Dosage and Administration:	This section contains file attachment.
----------------------------	--

General Directions:	NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER THE APPROPRIATE LEGISLATION.
---------------------	---

Withholding Periods:	<p>MEAT: DO NOT USE less than 42 days before slaughter for human consumption.</p> <p>MILK: ZERO (0) days.</p>
----------------------	---

Trade Advice:	EXPORT SLAUGHTER INTERVAL (ESI): DO NOT USE less than 42 days before slaughter for export. The ESI on this label was correct at the time of label approval. Before using this product, confirm the current ESI from Chemvet Australia Pty Ltd on 1800 243 683 or the APVMA website.
---------------	---

Safety Directions:	Harmful if swallowed. Will irritate the eyes and skin. Avoid contact with eyes and skin. When using the product wear cotton overalls buttoned to the neck and wrist (or equivalent clothing), elbow-length chemical resistant gloves and goggles. Wash hands after use. After each day's use, wash gloves, goggles and contaminated clothing
--------------------	--

First Aid Instructions:	<p>If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.</p> <p>If skin contact occurs, remove contaminated clothing and wash skin thoroughly. If in eyes, hold eyes open, flood with water for at least 15 minutes and see a doctor</p>
-------------------------	---

First Aid Warnings:	
---------------------	--

Additional User Safety:	
-------------------------	--

Environmental Statements:	<p><b>ENVIRONMENTAL PROTECTION:</b></p> <p>Ivermectin is extremely toxic to aquatic species. Do not contaminate dams, rivers, streams or other waterways with the chemical or used containers.</p>
---------------------------	--

Disposal:	<p><b>DISPOSAL (500mL, 1L)</b></p> <p>Dispose of empty container by wrapping with paper and putting in garbage.</p> <p><b>DISPOSAL (2.5L, 10L, 12.5L, 15L, 20L, 25L)</b></p> <p>Triple rinse containers before disposal then dispose of rinsings in a disposal pit away from desirable plants and watercourses. The disposal of undiluted chemicals is unacceptable. Destroy empty containers by breaking, crushing or puncturing or shredding them when applicable. Dispose of the containers at a local authority landfill readily available in your area. If no landfill is available, bury the containers at a depth of 500mm or more at a licence / approved disposable site. In some states wastes can only be buried at a licenced landfill. Do NOT burn empty containers or product. Clean up spillages promptly.</p>
-----------	---

Storage:	Store below 30°C (Room Temperature). Protect from light.
----------	--

## DOSAGE & ADMINISTRATION

VETMEC POUR-ON FOR CATTLE is for external application to cattle only. The recommended dose rate for VETMEC POUR-ON FOR CATTLE is 1mL/20kg body weight. Apply along the topline of the animal.

### DOSE RATE TABLE:

Live weight	Dose volume (mL)	500mL pack	1Litre pack	2.5Litre pack	10Litre pack	12.5Litre pack	15Litre pack	20Litre pack	25Litre pack
51-100 kg	5	100	200	500	2000	2500	3000	4000	5000
101-150 kg	7.5	66	133	333	1333	1666	2000	2666	3333
151-200 kg	10	50	100	250	1000	1250	1500	2000	2500
201-250 kg	12.5	40	80	200	800	1000	1200	1600	2000
251-300 kg	15	33	66	166	666	833	1000	1333	1666
301-350 kg	17.5	28	57	142	571	714	857	1142	1428
351-400 kg	20	25	50	125	500	625	750	1000	1250
401-450 kg	22.5	22	44	111	444	555	666	888	1111
451-500 kg	25	20	40	100	400	500	600	800	1000
501-550 kg	27.5	18	36	90	363	454	545	727	909
551-600 kg	30	16	33	83	333	416	500	666	833
601-650 kg	32.5	15	30	76	307	384	461	615	769

**A representative sample of animals should be weighed before treatment either with scales or a weigh band.**

Dose the mob according to the heaviest animal by live weight in the group (cows, bulls, calves, heifers, etc.). Do not underdose. Where there is a large variation in size within the group, draft into two or more lines based on body weight, to avoid excessive overdosing. For animals over 650kg (for example mature bulls) dose at a rate of 1 mL/20kg bodyweight.

Cattle should not be treated with VETMEC POUR-ON FOR CATTLE when the hide or hair is wet since the efficacy of the product may be reduced.

Rainfall on the herd within 2 hours of application may result in reduced efficacy of the product.