

Label: VETMEC F BROAD SPECTRUM ANTIPARASITIC INJECTION FOR BEEF AND DAIRY CATTLE  
Version: APVMA 290911  
Pack Size: 500 mL.

(Text above line is not included in the Label)

[Immediate container – front panel]

**CAUTION**

KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING  
FOR ANIMAL TREATMENT ONLY



**VETMEC F  
BROAD SPECTRUM ANTIPARASITIC INJECTION FOR BEEF AND DAIRY CATTLE**

IVERMECTIN 10 g/L  
CLORSULON 100 g/L



FOR THE TREATMENT AND CONTROL OF IVERMECTIN AND CLORSULON SENSITIVE  
STRAINS OF INTERNAL AND EXTERNAL PARASITES OF BEEF AND DAIRY CATTLE,  
INCLUDING ADULT LIVER FLUKE.

500 mL

(Text below this line not included in label)

Label: VETMEC F BROAD SPECTRUM ANTIPARASITIC INJECTION FOR BEEF AND DAIRY CATTLE  
Version: APVMA 290911  
Pack Size: 500 mL.

(Text above line is not included in the Label)

[Immediate container - back panel]

---

**READ CARTON BEFORE USING THIS PRODUCT**

**DIRECTIONS FOR USE**

The recommended dose is 1 mL / 50 kg liveweight by **subcutaneous injection only**. This product is not to be used intravenously or intramuscularly.

Live weight	Dose	500mL treats
Up to 100kg	2mL	250
101-150kg	3mL	166
151-200kg	4mL	125
201-250kg	5mL	100
251-300kg	6mL	83
301-350kg	7mL	71
351-400kg	8mL	62
401-450kg	9mL	55
451-500kg	10mL	50
501-550kg	11mL	45
551-600kg	12mL	41
601-650kg	13mL	38

Dose the mob according to the heaviest animal by live weight in the group (cows, bulls, calves, heifers, etc.). Do not underdose.

A representative sample of animals should be weighed before treatment either with scales or weighband. Cattle heavier than 650 kg, should be dosed at a rate of 1 mL / 50 kg body weight.

Where there is a large variation in size within the group, draft into two or more lines based on bodyweight, to avoid excessive overdosing.

**NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

**WITHHOLDING PERIODS:**

**MEAT: DO NOT USE less than 28 days before slaughter for human consumption.**

**MILK: Nil**

**TRADE ADVICE: Export Slaughter Interval (ESI):**

**DO NOT USE less than 42 days before slaughter for export. The ESI on this label was correct at the time of label approval. Before using this product confirm the current ESI from Chemvet Australia on 1800 243683 or the APVMA website [www.apvma.gov.au/residues/index.php](http://www.apvma.gov.au/residues/index.php)**

---

(Text below this line not included in label)

Label: VETMEC F BROAD SPECTRUM ANTIPARASITIC INJECTION FOR BEEF AND DAIRY CATTLE  
Version: APVMA 290911  
Pack Size: 500 mL.

(Text above line is not included in the Label)

[Immediate container - back panel - continued]

---

#### **SAFETY TO CATTLE**

Studies have demonstrated a wide margin of safety. Pregnant and breeding cattle may be treated provided normal care is taken in handling.

#### **SAFETY DIRECTIONS**

Poisonous if swallowed. Avoid contact with eyes, skin and clothing.

#### **FIRST AID**

If poisoning occurs, contact a doctor or the Poisons Information Centre, phone 131126. For further information refer to the Material Safety Data Sheet (MSDS) available at [www.chemvet.com.au](http://www.chemvet.com.au)

#### **DISPOSAL**

Dispose of empty container by wrapping with paper and putting in garbage. Discarded needles/sharps should immediately be placed in a designated and appropriately labelled 'sharps' containers.

#### **PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND THE ENVIRONMENT**

Ivermectin is very toxic to aquatic species. Do not contaminate dams, rivers, streams or other waterways with the chemical or used container

#### **STORAGE**

Store below 30°C (Room Temperature). Protect from light.

**CHEMVET Australia Pty Ltd**  
1/8 Rocklea Drive  
Port Melbourne VIC 3207  
[www.chemvet.com.au](http://www.chemvet.com.au)

Batch: APVMA 66768/54352  
Exp.

---

(Text below this line not included in label)

Label: VETMEC F BROAD SPECTRUM ANTIPARASITIC INJECTION FOR BEEF AND DAIRY CATTLE  
Version: APVMA 290911  
Pack Size: 500 mL.

(Text above line is not included in the Label)

[Carton – front panel]

---

CAUTION  
KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING  
FOR ANIMAL TREATMENT ONLY

**VETMEC F  
BROAD SPECTRUM ANTIPARASITIC INJECTION FOR BEEF AND DAIRY CATTLE**

IVERMECTIN 10 g/L  
CLORSULON 100 g/L

FOR THE TREATMENT AND CONTROL OF IVERMECTIN AND CLORSULON SENSITIVE  
STRAINS OF INTERNAL AND EXTERNAL PARASITES OF BEEF AND DAIRY CATTLE,  
INCLUDING ADULT LIVER FLUKE.

500 mL

CHEMVET AUSTRALIA Pty Ltd

---

(Text below this line not included in label)

Label: VETMEC F BROAD SPECTRUM ANTIPARASITIC INJECTION FOR BEEF AND DAIRY CATTLE  
Version: APVMA 290911  
Pack Size: 500 mL.

(Text above line is not included in the Label)

[Carton – back panel]

---

**VETMEC F BROAD SPECTRUM ANTIPARASITIC INJECTION FOR BEEF AND DAIRY CATTLE** contains ivermectin, a member of the macrocyclic lactone family of chemicals and clorsulon, a member of the sulphonamide family of chemicals.

When used at the recommended dose level of 200 µg ivermectin/kg and 2 mg clorsulon/kg liveweight, it is effective against sensitive strains of the following internal parasites:

**GASTRO-INTESTINAL ROUNDWORMS:**

Adult and immature:

*Ostertagia ostertagi* - Small brown stomach worm (including third and fourth stage larvae and inhibited immatures)

*Haemonchus placei* - Barber's pole worm

*Trichostrongylus axei* - Stomach hair worm (including third and fourth stage larvae)

*Trichostrongylus colubriformis* - Intestinal hair worm

*O. lyrata* - Stomach worm

*Cooperia spp* - Small intestinal worm (including third and fourth stage larvae)

*Oesophagostomum radiatum* - Nodule worm (including third and fourth stage larvae)

*Bunostomum phlebotomum* - Hookworm (including third and fourth stage larvae)

And adult stages of:

*Trichuris spp.* - Whipworm

*Strongyloides papillosus* - Intestinal worm

*Nematodirus spp.* - Thin necked intestinal worm

**LUNGWORMS**

*Dictyocaulus viviparus* (adult and immature, including inhibited stages)

**ADULT LIVER FLUKE**

*Fasciola hepatica*

**EYEWORM**

*Thelazia spp.* (adult stages)

**SUCKING LICE**

*Linognathus vituli* – Long nosed sucking louse

*Haematopinus eurysternus* – Short nosed sucking louse

*Solenopotes capillatus* – Little blue sucking louse

**MITES**

*Sarcoptes scabiei* var. *bovis*

**SCREW WORM FLY**

*Chrysomya bezziana* (Parasitic larval stages. Not reported in Australia, but present in Papua New Guinea)

---

(Text below this line not included in label)

Label: VETMEC F BROAD SPECTRUM ANTIPARASITIC INJECTION FOR BEEF AND DAIRY CATTLE  
Version: APVMA 290911  
Pack Size: 500 mL.

(Text above line is not included in the Label)

[Carton – back panel Continued]

---

## CATTLE TICK

*Boophilus microplus* – Cattle tick

When used as directed for the control of internal parasites of cattle, VETMEC F BROAD SPECTRUM ANTIPARASITIC INJECTION FOR BEEF AND DAIRY CATTLE concurrently aids in the control of:

## MITES

*Chorioptes bovis*

## BITING LICE

*Bovicola (Damalinia) bovis*

## SUSTAINED ACTIVITY:

At the recommended dose rate VETMEC F BROAD SPECTRUM ANTIPARASITIC INJECTION FOR BEEF AND DAIRY CATTLE effectively controls infection with *Ostertagia* spp, and *Cooperia* spp acquired up to 7 days after treatment and *Dictyocaulus viviparus* and *Chrysomyia bezziana* acquired up to 14 days after treatment.

## DIRECTIONS FOR USE

**DOSE RATE:** The recommended dose is 1 mL / 50 kg liveweight (200 µg ivermectin/kg and 2 mg clorsulon/kg liveweight) by **subcutaneous injection**. This product is not to be used intravenously or intramuscularly.

Liveweight	Dose	500 mL treats
Up to 100kg	2mL	250
101-150kg	3mL	166
151-200kg	4mL	125
201-250kg	5mL	100
251-300kg	6mL	83
301-350kg	7mL	71

### Caution: Avoid carcass damage

1. Sterilise all injection apparatus by boiling (or equivalent) before use. Avoid use of strong disinfectants on apparatus.
2. Maintain cleanliness at all times.
3. Keep needles sharp and clean. Replace frequently.
4. Use shortest needle possible, certainly not exceeding 15mm.
5. As far as possible avoid injection of animals during wet weather or under dusty conditions.
6. This product must be injected

---

(Text below this line not included in label)

Label: VETMEC F BROAD SPECTRUM ANTIPARASITIC INJECTION FOR BEEF AND DAIRY CATTLE  
Version: APVMA 290911  
Pack Size: 500 mL.

(Text above line is not included in the Label)

[Carton – back panel Continued]

---

351-400kg	8mL	62	only under the skin.  7. If possible inject high on the neck behind the ear.
401-450kg	9mL	55	
451-500kg	10mL	50	
501-550kg	11mL	45	
551-600kg	12mL	41	
601-650kg	13mL	38	

Dose the mob according to the heaviest animal by liveweight in each group (cows, bulls, calves, heifers, etc.). Do not underdose. A representative sample of cattle should be weighed before treatment either with scales or weighband. Cattle heavier than 650 kg, dose at rate of 1 mL / 50 kg body weight. Where there is a large variation in size within the group, draft into two or more lines based on bodyweight, to avoid excessive overdosing.

Resistance may develop to any drench

Consult your local Veterinary Advisor for recommended parasite management practices in your area

---

(Text below this line not included in label)

Label: VETMEC F BROAD SPECTRUM ANTIPARASITIC INJECTION FOR BEEF AND DAIRY CATTLE  
Version: APVMA 290911  
Pack Size: 500 mL.

(Text above line is not included in the Label)

[Carton – Side panel 1]

---

**NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

**WITHHOLDING PERIODS:**

**MEAT: DO NOT USE LESS THAN 28 DAYS BEFORE SLAUGHTER FOR HUMAN CONSUMPTION.**

**MILK: NIL**

**TRADE ADVICE: Export Slaughter Interval (ESI):**

**DO NOT USE less than 42 days before slaughter for export. The ESI on this label was correct at the time of label approval. Before using this product confirm the current ESI from Chemvet Australia on 1800 243683 or the APVMA website [www.apvma.gov.au/residues/index.php](http://www.apvma.gov.au/residues/index.php)**

**SAFETY TO CATTLE**

**Studies have demonstrated a wide margin of safety. Pregnant and breeding cattle may be treated provided normal care is taken in handling.**

**SAFETY DIRECTIONS**

**POISONOUS IF SWALLOWED. AVOID CONTACT WITH EYES, SKIN AND CLOTHING.**

**FIRST AID**

**If poisoning occurs, contact a doctor or Poisons Information Centre Ph 13 11 26. For further information refer to the Material Safety Data Sheet (MSDS) available at [www.chemvet.com.au](http://www.chemvet.com.au)**

**DISPOSAL**

**Dispose of empty container by wrapping with paper and putting in garbage. Discarded needles/sharps should immediately be placed in a designated and appropriately labelled 'sharps' containers.**

**PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND THE ENVIRONMENT**

**Ivermectin is very toxic to aquatic species. Do not contaminate dams, rivers, streams or other waterways with the chemical or used container**

---

(Text below this line not included in label)



Label: VETMEC F BROAD SPECTRUM ANTIPARASITIC INJECTION FOR BEEF AND DAIRY CATTLE  
Version: APVMA 290911  
Pack Size: 500 mL.

(Text above line is not included in the Label)

[Carton – Side panel 2]

---

**STORAGE**

Store below 30°C (Room Temperature). Protect from light.

APVMA Approval No: 66768/54352

Batch No:

Batch Expiry:

CHEMVET Australia Pty Ltd  
1/8 Rocklea Drive  
Port Melbourne VIC 3207  
[www.chemvet.com.au](http://www.chemvet.com.au)

---

(Text below this line not included in label)